



Weed TERMINATOR 20

A Horticultural Vinegar Biopesticide

For Non-Selective Control of Herbaceous Broadleaf Weeds and Weed Grasses which Surround Food crops, Non-food crops and Non-production Agricultural, Farmstead, Right-of-Way, and Institutional Land Sites

Active Ingredient:	32 Fl. Oz. (1 Qt) (0.946 L)
Acetic Acid	20.0%*
Other Ingredients	80.0%
Total	100.0%
*Equivalent to 200 grain vinegar by titration	

See directions for use in attached booklet.

DANGER – PELIGRO KEEP OUT OF REACH OF CHILDREN

FIRST AID
IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.



Manufactured for: Contact Organics (USA) 4947 US 75 Avenue MAURICE, IOWA 51036
info@contactorganics.com • EPA Reg. No. 85208-1-93740 EPA Est. No. 93740-THA-1

PERSONAL PROTECTION EQUIPMENT (PPE):

Applicators and other handlers must wear appropriate protective eyewear, such as face shield or goggles and face mask (with MSHA/NIOSH approval number prefix such as N95, R-95, or P-95), long sleeved shirt and long pants, waterproof gloves and shoes plus socks.

USER SAFETY REQUIREMENTS:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for PPE, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

- Users should:**
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards:

- This pesticide is toxic to birds exposed to spray drift, direct treatment or residues on crops or weeds. Do not apply this product or allow to drift to crops or weeds if birds are actively visiting the treatment area.
- This pesticide is toxic to fish and aquatic invertebrates.
- For terrestrial use only. Do not apply directly to water.

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Avoid application to reactive metals such as aluminum, tin, iron or items such as fencing or lawn furniture in order to prevent staining, mottling, or otherwise interfering with finished metal surfaces. In case of contact, rinse the sprayed surfaces with water. Avoid

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Annual Broadleaf Weeds

- Black medic
- Chickweed
- Black Nighthshade
- Cinquefoil (silvery)
- Dandelion
- Ground ivy
- Plantain spp.
- Toadflax
- Tufted vetch
- Wild carrot
- Moss
- Lambsquarters
- Mustard spp.
- Oxalis spp.
- Pigweed spp.
- Ragweed spp.
- Shepherdspurse
- Smartweed (Ladysthumb)
- Velvetleaf

Perennial Broadleaf Weeds

- Amaranth, spp
- Cinquefoil (silvery)

Perennial Grasses and Sedge

- Bluegrass
- Quackgrass
- Witchgrass

Liverworts*

*For Liverworts, dilute this product at a rate of 1 gallon of this product to 2 gallons of water. For all other weeds, use full strength.

Annual Grasses

- Crabgrass
- Foxtail spp.
- Italian ryegrass
- Poa annua

DO NOT SPRAY DIRECTLY ONTO FOOD CROPS

Use this product to control weeds that surround all food crops, nonfood crops (ornamentals) and non-production agricultural, farmstead, right-of-way, and institutional land sites. Use to control weeds in interstate freeways/highways, federal and state highways and city/county roads and walkways, railroad rights-of-way, tank farms, power stations and easements, and other rights-of-way, industrial sites, and vacant lots. Suitable to keep down weeds on all botanical and private association ornamental gardens, kennels, dog runs, and other animal enclosures and walkways.

DRIFT MANAGEMENT

- Drift is the off-site movement of spray away from the intended target. Managing drift is important for the protection of endangered species. In addition, this product is non-selective so it will severely injure or kill any crop plants or other plants it comes in contact with.
- Do not apply this product by aerial application.

NOTE: ALL CONTACTED VEGETATION WILL BE AFFECTED. DO NOT ALLOW THIS PRODUCT TO COME IN CONTACT WITH DESIRABLE PLANTS. OVERSPRAY OR DRIFT WILL INJURE OR KILL CONTACTED VEGETATION INCLUDING ANY CROP PLANTS.

Turf and Ornamentals -

Such as but not limited to woody ornamentals, potted crops, foliage plants, greenhouses, shadehouses, and any other plant production structure in which the product can be applied without contacting desirable plants. Applications to weeds in turf will kill surrounding turf.

Crops -

For all crops listed below, do not broadcast within the growing crop. Apply only with shielded or hooded spray equipment to weeds growing between crop rows. Any contact with the crop will cause crop injury. For all tree crops and fruit crops listed below, spray must be directed under the crop canopy. In crops where cultivation is possible, cultivate crop middles and reduce the sprayed area by directing the application to the uncultivated base of crop plants.

Citrus fruit -

Such as but not limited to orange, grapefruit, lemon, lime.

Pome fruit -

Such as but not limited to apple and pear.

Stone fruit -

Such as but not limited to cherry, peach, nectarine, plum.

Tree nuts -

Such as but not limited to almond, pecan, pistachio, coconut, betel nut, cashew, walnut.

Tropical fruit -

Such as but not limited to avocado, guava, jujube, noni, mango, star fruit, sugar apple, papaya, lychee, passion fruit, banana, date, olive, tamarind, pineapple.

Berries and small fruit -

Such as but not limited to strawberry, blackberry, raspberry, blueberry, cranberry, grapes, lingonberry, Saskatoon berry, kiwi fruit, maypop.

Root and tuber vegetables and leaves of root and tuber vegetables -

Such as but not limited to carrot, potato, radish, sugar beet, turnip tops, garden beet leaves, sugar beet leaves.

Leafy vegetables -

Such as but not limited to celery, head lettuce, leaf lettuce, spinach.

Brassica vegetables (cole crops) -

Such as but not limited to broccoli, cauliflower, cabbage, collard, mustard greens.

Fruiting vegetables -

Such as but not limited to tomato, bell pepper, chili pepper, eggplant, okra.

Mushrooms -

Apply preplant only. Do not apply directly to mushroom beds.

FOR SPOT TREATMENT APPLICATION:

Direct spray with a single nozzle or a hand-gun applicator to thoroughly wet undesirable weed foliage. Re-treatment of perennial weed growth may be required for control.

FOR BROADCAST SPRAY APPLICATIONS:

For broadcast spray application to control weeds. Application types include pre-plant (prior to crop establishment), preharvest desiccation (to kill crop foliage prior to harvesting the crop) postharvest crop desiccation (remaining crop residues after crop has been harvested).

Use 4.4 to 8.8 fluid ounces of this product per 100 square feet.

One quart will treat approximately 360 to 720 square feet.

Thorough coverage is necessary to achieve desirable control.

For dense weed populations and/or heavy foliage, use the higher spray volume to ensure adequate coverage. Apply with flat fan nozzles and at the lowest pressure needed to achieve required coverage, but do not exceed 35 PSI. If treating along the side of crop plants, shielded or hooded sprayers must be used to avoid crop injury.

SPRAY EQUIPMENT:

- Do not apply this product by aerial application.
- Do not apply when wind conditions favor movement of spray away from the site of application.
- When applying this product in a broadcast application utilize a hooded sprayer, shielded, or other spray equipment that prevents the movement of spray away from the target. Utilize nozzles that are designed to produce large spray droplets to reduce drift.
- Spray nozzles need to be plastic or 316 stainless steel as this product will eventually corrode any other metal.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE:

Keep from freezing. Store only in original tightly sealed container and out of reach of children.

FOR CONTROL OF LARGER ANNUALS AND BURNDOWN OF PERENNIAL WEED GROWTH:

Larger annual weeds and perennials are more difficult to control and may require retreatment. Treat initially as recommended above and repeat if new growth of leaves appears. Thorough coverage of all weed foliage is necessary to achieve desirable control. To reduce the amount of product required to control well-established weed stands, mow to a height of 4-6 inches and remove cut vegetation before applying this product.

TIMING OF APPLICATIONS:

Best results are achieved from applications to actively growing young weeds. Weeds that are mature, dormant, or hardened due to moisture stress are more tolerant of herbicide treatments. For best results, spray on dry leaves at temperatures between 50 and 75 degrees Fahrenheit. Conditions of high evaporation rates (temperatures higher than 75 degrees Fahrenheit and low relative humidity) may result in reduced efficacy. Only contact vegetation will be affected.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable container. Do no reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

WARRANTY STATEMENT:

To the extent required by law, Contact Organics (USA) warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label only when used in accordance with the label directions. Contact Organics (USA) makes no other express or implied warranties either of merchantability or fitness for a particular use and shall not be liable for misuse or conditions beyond its control, such as handling or storage.



Weed TERMINATOR 20

A Horticultural Vinegar Biopesticide


For Non-Selective Control of Herbaceous Broadleaf Weeds and Weed Grasses which Surround Food crops, Non-food crops and Non-production Agricultural, Farmstead, Right-of-Way, and Institutional Land Sites

Active Ingredient:	32 Fl. Oz. (1 Qt) (0.946 L)
Acetic Acid	20.0%*
Other Ingredients	80.0%
Total	100.0%
*Equivalent to 200 grain vinegar by titration	

See directions for use in attached booklet.

DANGER – PELIGRO KEEP OUT OF REACH OF CHILDREN

FIRST AID
IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.



Manufactured for: Contact Organics (USA) 4947 US 75 Avenue MAURICE, IOWA 51036
info@contactorganics.com • EPA Reg. No. 85208-1-93740 EPA Est. No. 93740-THA-1